

**ETUDE D'EFFICACITE ET DE TOLERANCE DE MIGRASTICK DANS LE  
TRAITEMENT DE LA MIGRAINE ( Dr SCHWOB)**

AIM OF THE STUDY	Evaluation of effectiveness and tolerance of Migrastick in treatment of migraines
METHODOLOGY	<ul style="list-style-type: none"> <li>- 21 patients</li> <li>- Pre-inclusion period of 15 days</li> <li>- Study duration : 2 months</li> <li>- Consultations at Day -15, Day 0, Day 60</li> <li>- <i>Inclusion criteria</i> : migraines for at least one year and at least one per week, migraine defined according to the headache classification criteria published by the International Headache Society in 1988.</li> <li>- <i>Evaluation criteria</i> : number of migraines, evaluation of the effectiveness and the tolerance duration of the migraine attacks, intensity of the pain and associated symptoms – nausea, vomiting) by the patient and the doctor (Bad to excellent)</li> </ul>
TREATMENTS	Migrastick is used in cases of migraine attacks and can be also used in combination with one of the 2 traditional medications against migraine predefined in the study.
RESULTS	<p><i>Number of migraines</i> : one every 3 days before the study, during the study one every 5 days.</p> <p><i>Evaluation of the effectiveness by the patient and the doctor</i>  Patient : 62% excellent, 29% good, 4.5% mediocre, 4.5 % bad  Doctor : 71 % excellent, 19% good, 10 % mediocre, 0 % bad</p> <p><i>Evaluation of the tolerance by the patient and the doctor</i>  Patient : 76% excellent, 9.5 % good, 9.5% mediocre, 5 % bad  Doctor : 71 % excellent, 19% good, 4.5 % mediocre, 4.5 % bad</p>
CONCLUSION	This study confirmed the effectiveness and the very good tolerance of Migrastick in migraines.

